SUBJECT: GUIDELINES FOR DETERMINING INTERFACILITY LEVEL OF TRANSPORT

(EMT, PARAMEDIC, MICN) REFERENCE NO. 517.1

PATIENT CONDITION / TREATMENT	BLS	ALS	SCT*
Stable patient who requires no special care, may have the following:			
 Foley catheter Gastrostomy tube Medication patch Nasogastric (NG) tube Patient controlled (or automated) medication pump that requires no intervention from transporting personnel 	✓		
Oxygen by mask or cannula	✓		
Heparin or saline lock	✓		
Peripheral IV without additives, maintained at a preset rate (normal saline, lactated ringers or glucose solutions)	✓		
Pre-existing vascular access device or peripherally inserted central catheter (PICC) line capped or with fluids infusing at a preset rate	✓		
Stable patient who requires cardiac monitoring or may need paramedic level intervention		✓	
Peripheral IV with any medication listed in the Los Angeles County paramedic scope of practice, infused with or without an IV pump		✓	
Chest tube clamped or attached to a Heimlich valve or similar device		✓	
Stable or unstable patient who requires care outside the paramedic scope of practice			✓
Continuous ventilatory assistance via a ventilator required (For stable patients, consider RCP staffed CCT transport)			✓
Any intervention or therapy that is not within the Los Angeles County paramedic scope of practice including IV infusion of any drug to an unstable patient			✓
Chest tube to suction			✓
Central venous catheter with central venous pressure (CVP) monitoring device in place			✓
Arterial line in place			✓
Pulmonary artery line in place (e.g. Swan Ganz)			✓
Intracranial pressure (ICP) line in place			✓
Intra-aortic balloon pump			✓
Transvenous pacemaker			✓
Standing orders written by transferring facility physician or provider agency medical director			✓

^{*}Includes RN and/or RCP staffed SCT transports – Consult with SCT Coordinator/Medical Director for specific indications for RN and/or RCP staffing of SCT transports.

EFFECTIVE: 01-01-13 REVISED: 01-01-23 SUPERSEDES: 04-01-19