

**COUNTY OF LOS ANGELES – DEPARTMENT OF MENTAL HEALTH
PROGRAM SUPPORT BUREAU – MHSA IMPLEMENTATION AND OUTCOMES DIVISION**



WELLNESS • RECOVERY • RESILIENCE

**Prevention & Early Intervention: Seeking Safety
Countywide Aggregate Practice Outcomes Dashboard Report
Outcome Data Submission through August 18, 2014**

Participating Legal Entities Include:

ALCOTT CENTER FOR MENTAL HEALTH SERVICES	HERITAGE CLINIC	TELECARE CORPORATION
ALMA FAMILY SERVICES	HILLSIDES	TESSIE CLEVELAND COMM SERVICES
ALMANSOR CLINICAL SERVICES	HILLVIEW MHC, INC.	THE GUIDANCE CENTER
AMANECER COMMUNITY COUNSELING SRVC	IMCES, INC.	THE HELP GROUP CHILD AND FAMILY CENTER
AVIVA CENTER	INTERCOMMUNITY CHILD GUIDANCE CENTER	THE VILLAGE FAMILY SERVICES
BARBOUR AND FLOYD ASSOCIATES	JEWISH FAMILY SERVICE OF LA	TOBINWORLD
BRIDGES, INC.	LEROY HAYNES FOUNDATION, INC.	TRINITY YOUTH SERVICES
CALIFORNIA HISPANIC COMMISSION ALCOHOL & DRUG ABUSE	LOS ANGELES CHILD GUIDANCE	UNITED AMERICAN INDIAN INVOLVEMENT
CHILD AND FAMILY CENTER	MCKINLEY CHILDRENS CENTER	VIP MENTAL HEALTH CENTER, INC.
CHILD AND FAMILY GUIDANCE CENTER	MENTAL HEALTH AMERICA OF LA	VISTA DEL MAR CHILD AND FAMILY SERVICES
CHILDREN'S HOSPITAL OF LOS ANGELES	OPTIMIST YOUTH HOMES	L.A. COUNTY DMH
COUNSELING AND RESEARCH ASSC, INC.	PACIFIC ASIAN COUNSELING SERVICES	AUGUSTUS F HAWKINS FAMILY MHS
COUNSELING4KIDS	PACIFIC CLINICS	AMERICAN INDIAN COUNSELING CENTER
CRITTENTON SERVICES FOR CHILD AND FAMILY	PACIFIC LODGE YOUTH SERVICES	ARCADIA MHS
D VEAL FAMILY AND YOUTH SERVICES	PASADENA UNIFIED SCHOOL DISTRICT	COASTAL API FAMILY MHC
DAVID AND MARGARET HOME, INC.	PENNY LANE CENTERS	COMPTON FAMILY MHS
DIDI HIRSCH PSYCHIATRIC SERVICE	PHOENIX HOUSE OF LOS ANGELES, INC.	DOWNTOWN MHC
DIGNITY HEALTH DBA CAL HOSP MED CTR	ROSEMARY CHILDRENS SERVICE	EDMUND D. EDELMAN WESTSIDE MHC
DREW CHILD DEVELOPMENT CORPORATION	SAINT JOHNS HEALTH CENTER	HARBOR-UCLA MEDICAL CENTER
EISNER PEDIATRIC FAMILY CENTER	SAN FERNANDO VALLEY COMMUNITY MHC	JUVENILE JUSTICE TRANSITION AFTERCARE SVCS
EL CENTRO DE AMISTAD, INC.	SAN GABRIEL CHILDRENS CENTER, INC.	LONG BEACH API FAMILY MHC
EL CENTRO DEL PUEBLO, INC.	SHIELDS FOR FAMILIES	LONG BEACH CHILD & ADOLESCENT CLINIC
ENKI HEALTH AND RESEARCH SYSTEMS, INC.	SOCIAL MODEL RECOVERY SYSTEMS	LONG BEACH MHS ADULT CLINIC
ETTIE LEE HOMES, INC.	SOUTH CENTRAL HEALTH AND REHAB PROGRAM	NORTHEAST MENTAL HEALTH CENTER
EXCEPTIONAL CHILDREN'S FOUNDATION	SPECIAL SERVICE FOR GROUPS	RIO HONDO COMMUNITY MHC
FAMILIES FIRST, INC.	SPIRITT FAMILY SERVICES	SAN PEDRO MENTAL HEALTH SERVICES
FIVE ACRES	ST. ANNES	SOUTH BAY MHS
FOOTHILL FAMILY SERVICES	ST. JOSEPH CENTER	WEST CENTRAL FAMILY MHS
HATHAWAY SYCAMORES CHILD FAMILY SERVICES	STIRLING ACADEMY, INC.	
HELPLINE YOUTH COUNSELING, INC.	TARZANA TREATMENT CENTERS, INC.	

Agencies submitting outcomes that are not approved to provide Seeking Safety by PEI Administration:

ASIAN AMERICAN DRUG ABUSE PROGRAM	ANTELOPE VALLEY MHS
VENICE FAMILY CLINIC	HOLLYWOOD MHC
L.A. COUNTY DMH	

# of Clients Claimed to Practice	# of Clients entered into PEI OMA	# of Tx cycles in PEI OMA	Clients with Multiple Tx Cycles	Clients Completing Tx	Clients Dropping-Out of Tx
22198	42.03%	9938	5.98%	23.49%	35.79%
n=	9329	n=	558	2334	3557

Note 1: Clients claimed was based on Seeking Safety being selected as the EBP in a PEI Plan and having ≥ 1 core services claimed to the practice.

Note 2: Number of clients Completing Tx or Dropping-Out of Tx was determined by whether the EBP was said to be completed (e.g. answered “yes” or “no”) in the PEI OMA.

Total Number of Clients	Age	Gender			Ethnicity					Primary Language		
	Average	Female	Male	Unknown	African-American	Asian / Pacific Islander	Caucasian	Hispanic / Latino	Other	English	Spanish	Other
9329	22	49.42%	50.52%	0.06%	20.82%	2.57%	11.50%	59.72%	5.39%	76.47%	20.46%	3.07%
n=		4610	4713	6	1942	240	1073	5571	503	7134	1909	286

Note1: Age is calculated at the date of the first EBP.

Note2: Percentages may not total 100 due to missing data and/or rounding.

Total Treatment Cycles	Depressive Disorder NOS	Oppositional Defiant Disorder	Post-Traumatic Stress Disorder	Mood Disorder NOS	Dysthymic Disorder	Other
9938	10.12%	9.60%	9.50%	8.67%	6.52%	55.58%
n=	1006	954	944	862	648	5524

*See breakout of “Other” on the supplemental information page.

Table 4. Program Process Data - Clients Who Entered Seeking Safety

Outcome Measures Administered	Pre-Test with Scores	Post-test with Scores	Clients Who Completed both a Pre and Post Measure with Scores
UCLA PTSD-RI - Parent	34.24%	16.80%	6.74%
n=	2272	646	447
Ackn=	6635	3846	6635
UCLA PTSD-RI - Child/Adolescent	64.70%	30.11%	13.47%
n=	4591	1221	956
Ackn=	7096	4055	7096
UCLA PTSD-RI - Adult	63.12%	30.28%	10.67%
n=	1402	291	237
Ackn=	2221	961	2221
Youth Outcome Questionnaire - 2.01 (Parent)	46.01%	22.33%	9.31%
n=	2916	816	590
Ackn=	6338	3655	6338
Youth Outcome Questionnaire – Self Report – 2.0 (YOQ-SR)	74.92%	37.02%	16.76%
n=	5081	1418	1137
Ackn=	6782	3830	6782
Outcome Questionnaire - 45.2	75.91%	38.21%	12.66%
n=	2033	426	339
Ackn=	2678	1115	2678

Note 1: Number of acknowledged measures (Ackn=) is determined by the number of required measures that receive a score or an unable to collect reason code.

Note 2: The % indicated for Pre-test with scores, Post-test with scores, and both a Pre- and Post-test with scores is calculated by dividing the (n=#) by the number acknowledged (Ackn=#) in the PEI OMA system for each measure. The number acknowledged (Ackn=#) for those with Pre and Post scores is an estimate based on the greatest number of matches that could be expected given the number of Pre scores acknowledged.

Table 5a. Top Reasons Given for "Unable to Collect"							
UCLA PTSD-RI - Parent	Total Pre 4364	Parent/care provider unavailable	Administration date exceeds acceptable range	Outcome measure unavailable	Parent/care provider refused	Clinician not trained in outcome measure	Other Reasons
	Percent	57.08%	13.82%	9.23%	5.80%	5.34%	8.73%
	n	2491	603	403	253	233	381
	Total Post 3200	Parent/care provider unavailable	Premature termination	Lost contact with parent/care provider	Parent/care provider refused	Outcome measure unavailable	Other Reasons
	Percent	52.09%	19.94%	6.38%	5.56%	5.06%	10.97%
	n	1667	638	204	178	162	351

Table 5b. Top Reasons Given for "Unable to Collect"							
UCLA PTSD-RI - Child/Adolescent	Total Pre 2505	Administration date exceeds acceptable range	Client refused	Outcome measure unavailable	Client unavailable	Clinician not trained in outcome measure	Other Reasons
	Percent	27.23%	20.92%	16.09%	13.77%	10.26%	11.74%
	n	682	524	403	345	257	294
	Total Post 2834	Client unavailable	Premature termination	Client refused	Lost contact with client	Administration date exceeds acceptable range	Other Reasons
	Percent	29.18%	28.44%	12.24%	10.66%	7.66%	11.82%
	n	827	806	347	302	217	335

Table 5c. Top Reasons Given for "Unable to Collect"							
UCLA PTSD-RI - Adult	Total Pre 819	Not available in primary language	Outcome measure unavailable	Clinician not trained in outcome measure	Client refused	Administration date exceeds acceptable range	Other Reasons
	Percent	35.78%	18.19%	17.46%	8.67%	8.42%	11.48%
	n	293	149	143	71	69	94
	Total Post 670	Lost contact with client	Client unavailable	Premature termination	Not available in primary language	Client refused	Other Reasons
	Percent	20.15%	19.40%	17.91%	17.31%	13.13%	12.09%
	n	135	130	120	116	88	81

Table 5d. Top Reasons Given for "Unable to Collect"							
Youth Outcome Questionnaire - 2.01 (Parent)	Total Pre 3422	Parent/care provider unavailable	Administration date exceeds acceptable range	Outcome measure unavailable	Parent/care provider refused	Clinician not trained in outcome measure	Other Reasons
	Percent	60.96%	16.66%	6.95%	5.46%	4.00%	5.96%
	n	2086	570	238	187	137	204
	Total Post 2839	Parent/care provider unavailable	Premature termination	Lost contact with parent/care provider	Parent/care provider refused	Administration date exceeds acceptable range	Other Reasons
	Percent	54.14%	21.52%	6.55%	5.42%	5.32%	7.04%
	n	1537	611	186	154	151	200

Table 5e. Top Reasons Given for "Unable to Collect"							
Youth Outcome Questionnaire Self Report – 2.0 (YOQ-SR)	Total Pre 1701	Administration date exceeds acceptable range	Client refused	Outcome measure unavailable	Client unavailable	Clinician not trained in outcome measure	Other Reasons
	Percent	35.21%	21.63%	15.70%	9.99%	8.41%	9.05%
	n	599	368	267	170	143	154
	Total Post 2412	Premature termination	Client unavailable	Lost contact with client	Client refused	Administration date exceeds acceptable range	Other Reasons
	Percent	31.76%	30.85%	11.28%	9.87%	7.34%	8.91%
	n	766	744	272	238	177	215

Table 5f. Top Reasons Given for "Unable to Collect"							
Outcome Questionnaire – 45.2	Total Pre 645	Outcome measure unavailable	Not available in primary language	Administration date exceeds acceptable range	Clinician not trained in outcome measure	Client refused	Other Reasons
	Percent	25.58%	17.36%	15.50%	14.11%	9.61%	17.83%
	n	165	112	100	91	62	115
	Total Post 689	Client unavailable	Lost contact with client	Premature termination	Client refused	Not available in primary language	Other Reasons
	Percent	25.11%	22.93%	22.50%	13.79%	5.52%	10.16%
	n	173	158	155	95	38	70

Table 6. Service Delivery Data – Clients Who Completed Seeking Safety						
Total Treatment Cycles 2334	Average Length of Treatment in Weeks	Range of Treatment Weeks		Average Number of Sessions	Range of Sessions	
	31	Min 0	Max 150	35	Min 1	Max 334

Table 7. Outcome Data* – Clients who Completed Seeking Safety

		Percent Improvement from Pre to Post	Percent of Clients Showing Reliable Change* from Pre-SS to Post-SS		
			Positive Change	No change	Negative Change
UCLA PTSD-RI - Parent	TOTAL	33.91%	23.38%	71.43%	5.19%
		(n=308)	72	220	16
UCLA PTSD-RI - Child/Adolescent	TOTAL	28.84%	26.21%	68.09%	5.70%
		(n=702)	184	478	40
UCLA PTSD-RI - Adult	TOTAL	25.84%	29.84%	67.02%	3.14%
		(n=191)	57	128	6
Youth Outcome Questionnaire (YOQ) - 2.01 (Parent)	TOTAL	33.30%	54.27%	32.66%	13.07%
		(n=398)	216	130	52
Youth Outcome Questionnaire - Self Report - 2.0 (YOQ-SR)	TOTAL	32.32%	45.51%	43.89%	10.60%
		(n=868)	395	381	92
Outcome Questionnaire – 45.2	TOTAL	25.63%	48.57%	45.36%	6.07%
		(n=280)	136	127	17

*Please see Appendix A. for a description of the Seeking Safety outcome measures and the outcome indicators (percent improvement in average scores; and, percent of clients showing reliable change).

Note1: Possible PTSD-RI range from 0-68, with a clinical cutpoint of 38.



Note2: Possible YOQ Total Scores range from -16-240, with a clinical cutpoint of 46.

Note3: Possible YOQ-SR Total Scores range from -16-240, with a clinical cutpoint of 47.



Note4: Possible OQ Total Scores range from 0-180, with a clinical cutpoint of 64.

Note5: Positive Change indicates that the scores decreased from the pre to the post measures.



UCLA PTSD RI – Parent
(n=308)





UCLA PTSD RI – Child/Adolescent
(n=702)





UCLA PTSD-RI – Adult (n=191)





Youth Outcome Questionnaire (YOQ) – 2.01 (Parent)
(N=398)



Youth Outcome Questionnaire – Self Report – 2.0 (YOQ-SR)
(N=868)



Outcome Questionnaire-45.2
(N=280)



Appendix

Youth Outcomes Questionnaires (YOQ and YOQ-SR)

The Youth Outcome Questionnaire is a 64-item parent-report that assesses global distress in a child's/adolescent's life from 4-17 years of age. The YOQ-SR is the Self-report version of the YOQ and is completed by the child/adolescent. Total scores on both measures can range from -16 to 240. Total scores of 46 or higher are most similar to a clinical population on the YOQ. A total score of 47 is most similar to that of a clinical population on the YOQ-SR.

Outcomes Questionnaires (OQ)

The Outcome Questionnaire is a 45-item self-report that assesses global distress in a client's life from ages 19 and older. Total Scores on this measure can range from 0 to 180, with scores of 64 or higher indicating clinical significance.

Post-Traumatic Stress Disorder Reaction Index (PTSD-RI)

The UCLA Post-Traumatic Stress Disorder Reaction Index (PTSD-RI) is a brief (21 or 22-item measure depending on the version) that measures the frequency of occurrence of post-traumatic stress disorder symptoms during the month prior to the assessment. The Child/Adolescent Version is appropriate for clients age 6-20. The Parent Version is appropriate for clients age 3-18. The Adult Short-form is appropriate for clients age 21+.

Possible Total PTSD Severity Scores range from 0-68; and scores of 38 or higher have the greatest sensitivity and specificity for detecting PTSD.

Reliable Change Index (RCI)

When comparing Pre and Post scores, it is very helpful to know whether the change reported represents the real effects of the treatment or errors in the system of measurement. The Reliability of Change Index (RCI) is a statistical way of helping to insure that the change recorded between pre and post assessments exceeds that which would be expected on the basis of measurement error alone. The RCI has been calculated using the Jacobson and Truax (1991) method and indicates when change exceeds that which would be expected on the basis of error at the $p < .05$ probability level. For a more in-depth discussion of Reliability of Change see Jacobson, N. S., & Truax, P. (1991). Clinical Significance: A statistical approach to defining meaningful change in psychotherapy research. *Journal of Consulting and Clinical Psychology*, 59, 12-19. Also see Wise, E. A. (2004). Methods for analyzing psychotherapy outcomes: A review of clinical significance, reliable change, and recommendations for future directions. *Journal of Personality Assessment*, 82(1), 50-59.

The number and percent of clients experiencing positive change, no change and negative change are recorded in table 7. Healthful change in each of the measures cited here means that scores have decreased in value from pre to post test administrations (i.e. recorded a negative change on the RCI). To help avoid confusion, healthful reliable change is presented as positive while unhealthy reliable change is presented as negative change.

The Reliable Change analysis for the PTSD-RI Adult Short-Form used the same Cronbach Alpha score found in the other PTSD-RI versions.

Supplemental Information for Table 3 “Other” Diagnosis Category

DSM-IV Primary Axis Diagnosis - Clients Who Entered Seeking Safety	n	Percent
Disruptive Behavior Disorder NOS	632	11.44%
Major Depressive Disorder, Recurrent, Moderate	479	8.67%
Conduct Disorder, Adolescent-Onset Type	457	8.27%
Attention-Deficit/Hyperactivity Disorder, Combined Type or Hyperactive Impulse Type	352	6.37%
Major Depressive Disorder, Rec., Severe W/O Psychotic Features	343	6.21%
Anxiety Disorder NOS	253	4.58%
Generalized Anxiety Disorder	239	4.33%
Major Depressive Disorder, Single Episode, Moderate	221	4.00%
Bipolar Disorder NOS	211	3.82%
Major Depressive Disorder, Rec., Severe With Psychotic Features	171	3.10%
Major Depressive Disorder, Recurrent, Unspecified	141	2.55%
Adjustment Disorder With Anxiety and Depressed Mood	120	2.17%
Major Depressive Disorder, SE, Severe W/O Psychotic Features	106	1.92%
Adjustment Disorder With Depressed Mood	106	1.92%
Major Depressive Disorder, Single Episode, Unspecified	106	1.92%
Psychotic Disorder NOS	106	1.92%
Major Depressive Disorder, Recurrent, Mild	101	1.83%
Major Depressive Disorder, Single Episode, Mild	91	1.65%
Schizoaffective Disorder	88	1.59%
Conduct Disorder, Childhood-Onset Type	82	1.48%
Attention-Deficit/Hyperactivity Disorder, Predominantly Inattentive Type	81	1.47%
Adjustment Disorder W/Mixed Disturbance Emotion and Conduct	76	1.38%
Major Depressive Disorder, SE, Severe With Psychotic Features	74	1.34%
Schizophrenia, Paranoid Type	62	1.12%
Attention-Deficit/Hyperactivity Disorder NOS	47	0.85%
Bipolar II Disorder	46	0.83%
Adjustment Disorder With Anxiety	44	0.80%
Panic Disorder Without Agoraphobia	43	0.78%
Adjustment Disorder Unspecified	40	0.72%
Adjustment Disorder With Disturbance of Conduct	32	0.58%
Bipolar I Disorder, MRE, Depressed, Severe W/O Psychotic Features	28	0.51%
Bipolar I Disorder, MRE, Depressed, Unspecified	25	0.45%
Bipolar I Disorder, MRE, Depressed, Severe With Psychotic Features	25	0.45%
Conduct Disorder, Unspecified Onset	25	0.45%
Intermittent Explosive Disorder	25	0.45%
Bipolar I Disorder, MRE, Mixed, Unspecified	21	0.38%
Bipolar I Disorder, MRE, Mixed, Moderate	20	0.36%
Obsessive-Compulsive Disorder	19	0.34%
Diagnosis Deferred (code invalid as Secondary/Axis II Diagnosis)	19	0.34%

DSM-IV Primary Axis Diagnosis - Clients Who Entered Seeking Safety	n	Percent
Bipolar I Disorder, MRE, Mixed, Severe W/O Psychotic Features	19	0.34%
Social Phobia	18	0.33%
Major Depressive Disorder, Recurrent, In Partial Remission	17	0.31%
Bipolar I Disorder, MRE, Depressed, Moderate	16	0.29%
Bipolar I Disorder, MRE, Mixed, Severe With Psychotic Features	16	0.29%
Panic Disorder With Agoraphobia	16	0.29%
No Diagnosis (code invalid as Secondary/Axis III Diagnosis)	15	0.27%
Bipolar I Disorder, MRE, Manic, Severe With Psychotic Features	15	0.27%
Bipolar I Disorder, MRE, Manic, Moderate	13	0.24%
Mood Disorder Due to ...(Indicate Medical Condition)	13	0.24%
Bipolar I Disorder, MRE, Unspecified	12	0.22%
Bipolar I Disorder, MRE, Manic, Severe W/O Psychotic Features	12	0.22%
Bipolar I Disorder, MRE, Hypomanic/Manic Unspecified	12	0.22%
Bipolar I Disorder, MRE, Depressed, Mild	11	0.20%
Acute Stress Disorder	10	0.18%
Cyclothymic Disorder	8	0.14%
Schizophrenia, Undifferentiated Type	8	0.14%
Schizophrenia, Disorganized Type	8	0.14%
Major Depressive Disorder, Single Episode, In Partial Remission	7	0.13%
Separation Anxiety Disorder	7	0.13%
Cannabis Abuse	7	0.13%
Reactive Attachment Disorder of Infancy or Early Childhood	7	0.13%
Pervasive Developmental Disorder NOS/Asperger's Disorder/Rett's Disorder	6	0.11%
Disorder of Infancy, Childhood, or Adolescence NOS	5	0.09%
Impulse-Control Disorder NOS	5	0.09%
Bipolar I Disorder, Single Manic Episode, Unspecified	5	0.09%
Bereavement	4	0.07%
Delusional Disorder	4	0.07%
Drug Induced Mood Disorder	4	0.07%
Bipolar I Disorder, MRE, Depressed, In Partial Remission	4	0.07%
Eating Disorder NOS	3	0.05%
Gender Identity Disorder in Adolescents or Adults	3	0.05%
Drug Induced Psychotic Disorder, With Delusions	3	0.05%
Drug Induced Psychotic Disorder, With Hallucinations	3	0.05%
Bipolar I Disorder, MRE, Mixed, Mild	3	0.05%
Alcohol-Induced Anxiety. Mood, Sleep Disorders/Sexual Dysfunction	3	0.05%
Anxiety Disorder Due to General Medical Condition	2	0.04%
Polysubstance Dependence	2	0.04%
Religious or Spiritual Problem/Phase of Life Problem	2	0.04%
Schizophrenia, Residual Type	2	0.04%
Major Depressive Disorder, Recurrent, In Full Remission	2	0.04%
Identity Problem	2	0.04%

DSM-IV Primary Axis Diagnosis - Clients Who Entered Seeking Safety	n	Percent
Bipolar I, SME, Severe W/O Psychotic Features	2	0.04%
Anorexia Nervosa	2	0.04%
Somatoform Disorder NOS and Undifferentiated Somatoform Disorder	2	0.04%
Bipolar I Disorder, Single Manic Episode, Moderate	2	0.04%
Dementia due to Medical Disorder/HIV Disease	1	0.02%
Dyspareunia (Not Due to a Medical Condition)	1	0.02%
Bulemia Nervosa	1	0.02%
Drug Induced Disorder NEC	1	0.02%
Schizophreniform Disorder	1	0.02%
Amphetamine Dependence	1	0.02%
Adult Antisocial Behavior	1	0.02%
Catatonic Disorder Due to General Medical Condition	1	0.02%
Cannabis Dependence	1	0.02%
Bipolar I Disorder, MRE, Manic, In Partial Remission	1	0.02%
Drug Related Disorder NOS	1	0.02%
Pain Disorder Associated w/Psychological and Medical	1	0.02%
Brief Psychotic Disorder	1	0.02%
Opioid Dependence	1	0.02%
Dissociative Identity Disorder	1	0.02%
Alcohol Dependence	1	0.02%
Unspecified Mental Disorder (Nonpsychotic)	1	0.02%
Major Depressive Disorder, Single Episode, In Full Remission	1	0.02%
Agoraphobia Without History of Panic Disorder	1	0.02%
Bipolar I Disorder, MRE Manic, In Full Remission	1	0.02%
Bipolar I, SME, Severe With Psychotic Features	1	0.02%
Psychotic Disorder Due to ...(Indicate Medical Condition) with Delusions	1	0.02%
Bipolar I Disorder, MRE, Depressed, In Full Remission	1	0.02%
Parent-Child Relational Problem	1	0.02%
Drug Induced Delirium	1	0.02%
TOTAL	5524	100.00%