

**COUNTY OF LOS ANGELES – DEPARTMENT OF MENTAL HEALTH
PROGRAM SUPPORT BUREAU – MHSA IMPLEMENTATION AND OUTCOMES DIVISION**



**Prevention & Early Intervention: Seeking Safety
Countywide Aggregate Practice Outcomes Dashboard Report
Outcome Data Submission through February 3, 2015**

Participating Legal Entities Include:

ALCOTT CENTER FOR MENTAL HEALTH SVC	HERITAGE CLINIC	TARZANA TREATMENT CENTERS INC
ALMA FAMILY SERVICES	HILLSIDES	TELECARE CORPORATION
AMANECER COMMUNITY COUNSELING SVC	HILLVIEW MENTAL HEALTH CENTER INC	TESSIE CLEVELAND COMM SRVC CORP
BARBOUR AND FLOYD MEDCL ASSOCIATES	IMCES INC	THE GUIDANCE CENTER
BRIDGES INC	INS FOR RDSGN OF LEARN DBA ALMANSOR	THE HELP GROUP CHILD AND FAMILY CTR
CA HISPANIC COMMISSION A D ABUSE	INTERCOMMUNITY CHILD GUIDANCE CTR	THE VILLAGE FAMILY SERVICES
CHILD AND FAMILY CENTER	JEWISH FAMILY SERVICE OF LOS ANGELES	TOBINWORLD
CHILD AND FAMILY GUIDANCE CENTER	LEROY HAYNES FOUNDATION INC	TRINITY YOUTH SERVICES
CHILDRENS HOSPITAL OF LOS ANGELES	LOS ANGELES CHILD GUIDANCE	UNITED AMERICAN INDIAN INVOLVEMENT
CHILDRENS INSTITUTE INC	MCKINLEY CHILDRENS CENTER	VIP COMMUNITY MENTAL HEALTH CTR INC
COUNSELING AND RESEARCH ASSC INC	MENTAL HEALTH AMERICA OF LA	VISTA DEL MAR CHILD AND FAMILY SVC
COUNSELING4KIDS	OLIVE CREST	L.A. COUNTY DMH
CRITTENTON SVS FOR CHILD AND FAMILIES	OPTIMIST YOUTH HOMES DBA BOYS HOME	EDMUND D. EDELMAN WESTSIDE
D VEAL FAMILY AND YOUTH SVC	PACIFIC ASIAN COUN SVC	AMERICAN INDIAN COUNSELING
DAVID AND MARGARET HOME INC	PACIFIC CLINICS	ARCADIA MHS
DIDI HIRSCH PSYCHIATRIC SVC	PACIFIC LODGE YOUTH SERVICES	AUGUSTUS F HAWKINS FAMILY
DIGNITY HEALTH DBA CAL HOSP MED CTR	PASADENA UNIFIED SCHOOL DISTRICT	COASTAL API FAMILY MHC
DREW CHILD DEVELOPMENT CORPORATION	PENNY LANE CENTERS	COMPTON FAMILY MHS
EISNER PEDIATRIC FAMILY CTR	PHOENIX HOUSE OF LOS ANGELES INC	DOWNTOWN MENTAL HEALTH
EL CENTRO DE AMISTAD INC	ROSEMARY CHILDRENS SERVICE	HARBOR-UCLA MEDICA
EL CENTRO DEL PUEBLO INC	SAINT JOHNS HEALTH CENTER	JUV JUS TRANSITION AFTERCARE
ENKI HEALTH AND RESEARCH SYSTEMS I	SAN FERNANDO VALLEY COMMUNITY MHC	LONG BEACH API FAMILY MHC
ETTIE LEE HOMES INC	SAN GABRIEL CHILDRENS CENTER INC	LONG BEACH CHILD ADOLESCENT CLINIC
EXCEPTIONAL CHILDRENS FOUNDATION	SHIELDS FOR FAMILIES	LONG BEACH MHS ADULT CLINIC
FAMILIES FIRST INC	SOCIAL MODEL RECOVERY SYSTEMS	NORTHEAST MENTAL HEALTH CENTER
FIVE ACRES	SOUTH CENTRAL HEALTH AND REHAB PROG	RIO HONDO COMMUNITY MHC
FOOTHILL FAMILY SERVICE	SPECIAL SERVICE FOR GROUPS	SAN PEDRO MENTAL HEALTH
HAMBURGER HOME DBA AVIVA CENTER	ST ANNES	SOUTH BAY MHS
HATHAWAY SYCAMORES CHILD FAM SVC	ST JOSEPH CENTER	TIES FOR FAMILIES
HELPLINE YOUTH COUNSELING INC	STIRLING ACADEMY INC	WEST CENTRAL FAMILY MHS

Agencies submitting outcomes that are not approved to provide Seeking Safety by PEI Administration:

ASIAN AMERICAN DRUG ABUSE PROGRAM	L.A. COUNTY DMH
VENICE FAMILY CLINIC	ANTELOPE VALLEY MHS
	HOLLYWOOD MHC

# of Clients Claimed to Practice	# of Clients entered into PEI OMA	# of Tx cycles in PEI OMA	Clients with Multiple Tx Cycles	Clients Completing Tx	Clients Dropping-Out of Tx	Clients Still in Tx
19824	56.65%	12026	6.38%	24.60%	38.58%	36.82%
n=	11230	n=	717	2958	4640	4428

Note 1: Clients claimed was based on Seeking Safety being selected as the EBP in a PEI Plan and having ≥ 2 core services claimed to the practice.

Note 2: Number of clients Completing Tx or Dropping-Out of Tx was determined by whether the EBP was said to be completed (e.g. answered “yes” or “no”) in the PEI OMA.

Total Number of Clients	Age	Gender			Ethnicity					Primary Language		
	Average	Female	Male	Unknown	African-American	Asian / Pacific Islander	Caucasian	Hispanic / Latino	Other	English	Spanish	Other
11230	22	50.32%	49.63%	0.05%	20.28%	2.40%	11.60%	59.88%	5.85%	76.61%	20.44%	2.96%
n=		5651	5573	6	2277	269	1303	6724	657	8603	2295	332

Note1: Age is calculated at the date of the first EBP.

Note2: Percentages may not total 100 due to missing data and/or rounding.

Total Treatment Cycles	Depressive Disorder NOS	Oppositional Defiant Disorder	Post-Traumatic Stress Disorder	Mood Disorder NOS	Disruptive Behavior Disorder NOS	Other
12026	10.35%	9.62%	9.36%	8.54%	6.53%	55.60%
n=	1245	1157	1126	1027	785	6686

*See breakout of “Other” on the supplemental information page.

Table 4. Program Process Data - Clients Who Entered Seeking Safety

Outcome Measures Administered	Pre-Test with Scores	Post-test with Scores	Clients Who Completed both a Pre and Post Measure with Scores
UCLA PTSD-RI - Parent	33.72%	16.33%	6.86%
n=	2740	802	557
Ackn=	8125	4912	8125
UCLA PTSD-RI - Child/Adolescent	63.75%	28.95%	13.67%
n=	5513	1497	1182
Ackn=	8648	5171	8648
UCLA PTSD-RI - Adult	62.22%	27.50%	11.44%
n=	1675	377	308
Ackn=	2692	1371	2692
Youth Outcome Questionnaire - 2.01 (Parent)	45.77%	21.93%	9.58%
n=	3527	1019	738
Ackn=	7706	4647	7706
Youth Outcome Questionnaire – Self Report – 2.0 (YOQ-SR)	74.43%	35.33%	16.90%
n=	6136	1725	1393
Ackn=	8244	4883	8244
Outcome Questionnaire - 45.2	76.29%	37.52%	15.00%
n=	2462	591	484
Ackn=	3227	1575	3227

Note 1: Number of acknowledged measures (Ackn=) is determined by the number of required measures that receive a score or an unable to collect reason code.

Note 2: The % indicated for Pre-test with scores, Post-test with scores, and both a Pre- and Post-test with scores is calculated by dividing the (n=#) by the number acknowledged (Ackn=#) in the PEI OMA system for each measure. The number acknowledged (Ackn=#) for those with Pre and Post scores is an estimate based on the greatest number of matches that could be expected given the number of Pre scores acknowledged.

Table 5a. Top Reasons Given for "Unable to Collect"

UCLA PTSD-RI - Parent	Total Pre 5386	Parent/care provider unavailable	Administration date exceeds acceptable range	Outcome measure unavailable	Parent/care provider refused	Clinician not trained in outcome measure	Other Reasons
	Percent	57.22%	14.46%	8.35%	6.78%	4.40%	8.78%
	n	3082	779	450	365	237	473
	Total Post 4110	Parent/care provider unavailable	Premature termination	Lost contact with parent/care provider	Parent/care provider refused	Administration date exceeds acceptable range	Other Reasons
	Percent	51.22%	20.19%	6.47%	5.72%	5.60%	10.80%
	n	2105	830	266	235	230	444

Table 5b. Top Reasons Given for "Unable to Collect"

UCLA PTSD-RI - Child/Adolescent	Total Pre 3135	Administration date exceeds acceptable range	Client refused	Outcome measure unavailable	Client unavailable	Clinician not trained in outcome measure	Other Reasons
	Percent	27.56%	23.19%	14.26%	14.10%	8.33%	12.57%
	n	864	727	447	442	261	394
	Total Post 3674	Premature termination	Client unavailable	Client refused	Lost contact with client	Administration date exceeds acceptable range	Other Reasons
	Percent	29.04%	28.36%	12.63%	10.42%	8.08%	11.46%
	n	1067	1042	464	383	297	421

Table 5c. Top Reasons Given for "Unable to Collect"

UCLA PTSD-RI - Adult	Total Pre 1017	Not available in primary language	Clinician not trained in outcome measure	Outcome measure unavailable	Client refused	Administration date exceeds acceptable range	Other Reasons
	Percent	34.22%	18.58%	16.81%	8.95%	8.85%	12.59%
	n	348	189	171	91	90	128
	Total Post 994	Not available in primary language	Client unavailable	Lost contact with client	Premature termination	Client refused	Other Reasons
	Percent	20.12%	19.52%	17.91%	16.80%	11.17%	14.49%
	n	200	194	178	167	111	144

Table 5d. Top Reasons Given for "Unable to Collect"

Youth Outcome Questionnaire - 2.01 (Parent)	Total Pre 4179	Parent/care provider unavailable	Administration date exceeds acceptable range	Parent/care provider refused	Outcome measure unavailable	Clinician not trained in outcome measure	Other Reasons
	Percent	60.90%	17.13%	6.29%	6.25%	3.37%	6.05%
	n	2545	716	263	261	141	253
	Total Post 3628	Parent/care provider unavailable	Premature termination	Lost contact with parent/care provider	Administration date exceeds acceptable range	Parent/care provider refused	Other Reasons
	Percent	53.03%	21.61%	6.62%	5.98%	5.57%	7.19%
	n	1924	784	240	217	202	261

Table 5e. Top Reasons Given for "Unable to Collect"

Youth Outcome Questionnaire Self Report - 2.0 (YOQ-SR)	Total Pre 2108	Administration date exceeds acceptable range	Client refused	Outcome measure unavailable	Client unavailable	Clinician not trained in outcome measure	Other Reasons
	Percent	35.44%	21.96%	13.95%	11.57%	7.02%	10.06%
	n	747	463	294	244	148	212
	Total Post 3158	Premature termination	Client unavailable	Lost contact with client	Client refused	Administration date exceeds acceptable range	Other Reasons
	Percent	32.33%	29.99%	10.80%	10.51%	7.79%	8.58%
	n	1021	947	341	332	246	271

Table 5f. Top Reasons Given for "Unable to Collect"

Outcome Questionnaire - 45.2	Total Pre 765	Outcome measure unavailable	Administration date exceeds acceptable range	Not available in primary language	Clinician not trained in outcome measure	Client refused	Other Reasons
	Percent	23.14%	17.12%	15.42%	12.29%	11.24%	20.78%
	n	177	131	118	94	86	159
	Total Post 984	Client unavailable	Lost contact with client	Premature termination	Client refused	Not available in primary language	Other Reasons
	Percent	25.51%	21.85%	21.65%	12.09%	7.42%	11.48%
	n	251	215	213	119	73	113

Table 6. Service Delivery Data – Clients Who Completed Seeking Safety						
Total Treatment Cycles 2958	Average Length of Treatment in Weeks	Range of Treatment Weeks		Average Number of Sessions	Range of Sessions	
	31	Min 0	Max 209	35	Min 1	Max 383

Table 7. Outcome Data* – Clients who Completed Seeking Safety

		Percent Improvement from Pre to Post	Percent of Clients Showing Reliable Change* from Pre-SS to Post-SS		
			Positive Change	No change	Negative Change
UCLA PTSD-RI - Parent	TOTAL	35.55%	26.17%	68.39%	5.44%
		(n=386)	101	264	21
UCLA PTSD-RI - Child/Adolescent	TOTAL	28.64%	25.71%	67.89%	6.40%
		(n=875)	225	594	56
UCLA PTSD-RI - Adult	TOTAL	25.56%	30.52%	66.27%	3.21%
		(n=249)	76	165	8
Youth Outcome Questionnaire (YOQ) - 2.01 (Parent)	TOTAL	35.34%	55.76%	32.53%	11.72%
		(n=495)	276	161	58
Youth Outcome Questionnaire - Self Report - 2.0 (YOQ-SR)	TOTAL	32.03%	43.71%	46.32%	9.97%
		(n=1073)	469	497	107
Outcome Questionnaire – 45.2	TOTAL	29.26%	53.02%	41.71%	5.28%
		(n=398)	211	166	21

*Please see Appendix A. for a description of the Seeking Safety outcome measures and the outcome indicators (percent improvement in average scores; and, percent of clients showing reliable change).

Note1: Possible PTSD-RI range from 0-68, with a clinical cutpoint of 38.



Note2: Possible YOQ Total Scores range from -16-240, with a clinical cutpoint of 46.

Note3: Possible YOQ-SR Total Scores range from -16-240, with a clinical cutpoint of 47.



Note4: Possible OQ Total Scores range from 0-180, with a clinical cutpoint of 64.

Note5: Positive Change indicates that the scores decreased from the pre to the post measures.



UCLA PTSD RI – Parent
(n=386)





UCLA PTSD RI – Child/Adolescent
(n=875)





UCLA PTSD-RI – Adult (n=249)





Youth Outcome Questionnaire (YOQ) – 2.01 (Parent)
(N=495)



Youth Outcome Questionnaire – Self Report – 2.0 (YOQ-SR)
(N=1073)



Outcome Questionnaire-45.2
(N=398)



Appendix

Youth Outcomes Questionnaires (YOQ and YOQ-SR)

The Youth Outcome Questionnaire is a 64-item parent-report that assesses global distress in a child's/adolescent's life from 4-17 years of age. The YOQ-SR is the Self-report version of the YOQ and is completed by the child/adolescent. Total scores on both measures can range from -16 to 240. Total scores of 46 or higher are most similar to a clinical population on the YOQ. A total score of 47 is most similar to that of a clinical population on the YOQ-SR.

Outcomes Questionnaires (OQ)

The Outcome Questionnaire is a 45-item self-report that assesses global distress in a client's life from ages 19 and older. Total Scores on this measure can range from 0 to 180, with scores of 64 or higher indicating clinical significance.

Post-Traumatic Stress Disorder Reaction Index (PTSD-RI)

The UCLA Post-Traumatic Stress Disorder Reaction Index (PTSD-RI) is a brief (21 or 22-item measure depending on the version) that measures the frequency of occurrence of post-traumatic stress disorder symptoms during the month prior to the assessment. The Child/Adolescent Version is appropriate for clients age 6-20. The Parent Version is appropriate for clients age 3-18. The Adult Short-form is appropriate for clients age 21+.

Possible Total PTSD Severity Scores range from 0-68; and scores of 38 or higher have the greatest sensitivity and specificity for detecting PTSD.

Reliable Change Index (RCI)

When comparing Pre and Post scores, it is very helpful to know whether the change reported represents the real effects of the treatment or errors in the system of measurement. The Reliability of Change Index (RCI) is a statistical way of helping to insure that the change recorded between pre and post assessments exceeds that which would be expected on the basis of measurement error alone. The RCI has been calculated using the Jacobson and Truax (1991) method and indicates when change exceeds that which would be expected on the basis of error at the $p < .05$ probability level. For a more in-depth discussion of Reliability of Change see Jacobson, N. S., & Truax, P. (1991). Clinical Significance: A statistical approach to defining meaningful change in psychotherapy research. *Journal of Consulting and Clinical Psychology*, 59, 12-19. Also see Wise, E. A. (2004). Methods for analyzing psychotherapy outcomes: A review of clinical significance, reliable change, and recommendations for future directions. *Journal of Personality Assessment*, 82(1), 50-59.

The number and percent of clients experiencing positive change, no change and negative change are recorded in table 7. Healthful change in each of the measures cited here means that scores have decreased in value from pre to post test administrations (i.e. recorded a negative change on the RCI). To help avoid confusion, healthful reliable change is presented as positive while unhealthful reliable change is presented as negative change.

The Reliable Change analysis for the PTSD-RI Adult Short-Form used the same Cronbach Alpha score found in the other PTSD-RI versions.

Supplemental Information for Table 3 “Other” Diagnosis Category

DSM-IV Primary Axis Diagnosis - Clients Who Entered Seeking Safety	n	Percent
Dysthymic Disorder	783	11.71%
Major Depressive Disorder, Recurrent, Moderate	598	8.94%
Conduct Disorder, Adolescent-Onset Type	540	8.08%
Attention-Deficit/Hyperactivity Disorder, Combined Type or Hyperactive Impulse Type	421	6.30%
Major Depressive Disorder, Rec., Severe W/O Psychotic Features	397	5.94%
Anxiety Disorder NOS	322	4.82%
Generalized Anxiety Disorder	300	4.49%
Major Depressive Disorder, Single Episode, Moderate	276	4.13%
Bipolar Disorder NOS	249	3.72%
Major Depressive Disorder, Rec., Severe With Psychotic Features	188	2.81%
Major Depressive Disorder, Recurrent, Unspecified	176	2.63%
Adjustment Disorder With Anxiety and Depressed Mood	149	2.23%
Major Depressive Disorder, Single Episode, Unspecified	136	2.03%
Adjustment Disorder With Depressed Mood	135	2.02%
Major Depressive Disorder, SE, Severe W/O Psychotic Features	127	1.90%
Major Depressive Disorder, Recurrent, Mild	123	1.84%
Psychotic Disorder NOS	121	1.81%
Major Depressive Disorder, Single Episode, Mild	118	1.76%
Attention-Deficit/Hyperactivity Disorder, Predominantly Inattentive Type	98	1.47%
Schizoaffective Disorder	97	1.45%
Adjustment Disorder W/Mixed Disturbance Emotion and Conduct	94	1.41%
Conduct Disorder, Childhood-Onset Type	93	1.39%
Major Depressive Disorder, SE, Severe With Psychotic Features	78	1.17%
Attention-Deficit/Hyperactivity Disorder NOS	67	1.00%
Schizophrenia, Paranoid Type	65	0.97%
Adjustment Disorder Unspecified	55	0.82%
Panic Disorder Without Agoraphobia	53	0.79%
Bipolar II Disorder	52	0.78%
Adjustment Disorder With Anxiety	48	0.72%
Adjustment Disorder With Disturbance of Conduct	42	0.63%
Bipolar I Disorder, MRE, Depressed, Severe W/O Psychotic Features	33	0.49%
Bipolar I Disorder, MRE, Depressed, Severe With Psychotic Features	33	0.49%
Bipolar I Disorder, MRE, Depressed, Unspecified	32	0.48%
Conduct Disorder, Unspecified Onset	31	0.46%
Intermittent Explosive Disorder	29	0.43%
Bipolar I Disorder, MRE, Mixed, Unspecified	24	0.36%
Panic Disorder With Agoraphobia	23	0.34%
Bipolar I Disorder, MRE, Mixed, Severe W/O Psychotic Features	22	0.33%
Obsessive-Compulsive Disorder	22	0.33%

DSM-IV Primary Axis Diagnosis - Clients Who Entered Seeking Safety	n	Percent
Social Phobia	22	0.33%
Bipolar I Disorder, MRE, Mixed, Moderate	22	0.33%
Diagnosis Deferred (code invalid as Secondary/Axis II Diagnosis)	22	0.33%
Bipolar I Disorder, MRE, Mixed, Severe With Psychotic Features	20	0.30%
Major Depressive Disorder, Recurrent, In Partial Remission	19	0.28%
Bipolar I Disorder, MRE, Depressed, Moderate	18	0.27%
Bipolar I Disorder, MRE, Manic, Severe With Psychotic Features	18	0.27%
No Diagnosis (code invalid as Secondary/Axis III Diagnosis)	16	0.24%
Mood Disorder Due to ...(Indicate Medical Condition)	14	0.21%
Bipolar I Disorder, MRE, Hypomanic/Manic Unspecified	14	0.21%
Bipolar I Disorder, MRE, Manic, Moderate	14	0.21%
Bipolar I Disorder, MRE, Unspecified	13	0.19%
Bipolar I Disorder, MRE, Depressed, Mild	13	0.19%
Bipolar I Disorder, MRE, Manic, Severe W/O Psychotic Features	12	0.18%
Acute Stress Disorder	12	0.18%
Schizophrenia, Undifferentiated Type	11	0.16%
Cyclothymic Disorder	11	0.16%
Reactive Attachment Disorder of Infancy or Early Childhood	10	0.15%
Schizophrenia, Disorganized Type	10	0.15%
Impulse-Control Disorder NOS	9	0.13%
Cannabis Abuse	9	0.13%
Major Depressive Disorder, Single Episode, In Partial Remission	8	0.12%
Separation Anxiety Disorder	8	0.12%
Pervasive Developmental Disorder NOS/Asperger's Disorder/Rett's Disorder	7	0.10%
Disorder of Infancy, Childhood, or Adolescence NOS	7	0.10%
Bipolar I Disorder, Single Manic Episode, Unspecified	6	0.09%
Major Depressive Disorder, Recurrent, In Full Remission	5	0.07%
Bipolar I Disorder, MRE, Depressed, In Partial Remission	4	0.06%
Drug Induced Mood Disorder	4	0.06%
Amphetamine Dependence	4	0.06%
Delusional Disorder	4	0.06%
Bereavement	4	0.06%
Eating Disorder NOS	4	0.06%
Gender Identity Disorder in Adolescents or Adults	3	0.04%
Drug Induced Psychotic Disorder, With Hallucinations	3	0.04%
Polysubstance Dependence	3	0.04%
Drug Induced Psychotic Disorder, With Delusions	3	0.04%
Bipolar I Disorder, MRE, Mixed, Mild	3	0.04%
Alcohol-Induced Anxiety. Mood, Sleep Disorders/Sexual Dysfunction	3	0.04%
Opioid Dependence	2	0.03%
Identity Problem	2	0.03%
Bipolar I Disorder, Single Manic Episode, Moderate	2	0.03%

DSM-IV Primary Axis Diagnosis - Clients Who Entered Seeking Safety	n	Percent
Anorexia Nervosa	2	0.03%
Bipolar I Disorder, MRE Manic, In Full Remission	2	0.03%
Major Depressive Disorder, Single Episode, In Full Remission	2	0.03%
Schizophrenia, Residual Type	2	0.03%
Religious or Spiritual Problem/Phase of Life Problem	2	0.03%
Somatoform Disorder NOS and Undifferentiated Somatoform Disorder	2	0.03%
Anxiety Disorder Due to General Medical Condition	2	0.03%
Bipolar I, SME, Severe W/O Psychotic Features	2	0.03%
Dissociative Identity Disorder	1	0.01%
Psychotic Disorder Due to ...(Indicate Medical Condition) with Delusions	1	0.01%
Drug Induced Disorder NEC	1	0.01%
Parent-Child Relational Problem	1	0.01%
Agoraphobia Without History of Panic Disorder	1	0.01%
Drug Related Disorder NOS	1	0.01%
Brief Psychotic Disorder	1	0.01%
Dementia due to Medical Disorder/HIV Disease	1	0.01%
Bipolar I Disorder, MRE, Depressed, In Full Remission	1	0.01%
Dyspareunia (Not Due to a Medical Condition)	1	0.01%
Adult Antisocial Behavior	1	0.01%
Catatonic Disorder Due to General Medical Condition	1	0.01%
Bipolar I Disorder, MRE, Manic, In Partial Remission	1	0.01%
Schizophreniform Disorder	1	0.01%
Alcohol Dependence	1	0.01%
Trichotillomania	1	0.01%
Bipolar I, SME, Severe With Psychotic Features	1	0.01%
Drug Induced Delirium	1	0.01%
Unspecified Mental Disorder (Nonpsychotic)	1	0.01%
Cannabis Dependence	1	0.01%
Bulimia Nervosa	1	0.01%
Pain Disorder Associated w/Psychological and Medical	1	0.01%
Total	6686	100.00%