

**COUNTY OF LOS ANGELES – DEPARTMENT OF MENTAL HEALTH
PROGRAM SUPPORT BUREAU – MHSA IMPLEMENTATION AND OUTCOMES DIVISION**



**Prevention & Early Intervention: Seeking Safety
Countywide Aggregate Practice Outcomes Dashboard Report
Outcome Data Submission through July 15, 2015**

Participating Legal Entities Include:

ALCOTT CENTER FOR MENTAL HEALTH SVC	HELPLINE YOUTH COUNSELING INC	STIRLING ACADEMY, INC.
ALMA FAMILY SERVICES	HERITAGE CLINIC	TARZANA TREATMENT CENTERS INC
AMANECER COMMUNITY COUNSELING SVC	HILLSIDES	TELECARE CORPORATION
BARBOUR AND FLOYD MEDCL ASSOCIATES	HILLVIEW MENTAL HEALTH CENTER INC	TESSIE CLEVELAND COMM SRVC CORP
BIENVENIDOS CHILDRENS CENTER INC	IMCES INC	THE GUIDANCE CENTER
BRIDGES INC	INS FOR RDSGN OF LEARN DBA ALMANSOR	THE HELP GROUP CHILD AND FAMILY CTR
CA HISPANIC COMMISSION A D ABUSE	INTERCOMMUNITY CHILD GUIDANCE CTR	THE VILLAGE FAMILY SERVICES
CHILD AND FAMILY CENTER	JEWISH FAMILY SERVICE OF LOS ANGELES	TOBINWORLD
CHILD AND FAMILY GUIDANCE CENTER	LEROY HAYNES FOUNDATION INC	TRINITY YOUTH SERVICES
CHILDRENS CENTER OF ANTELOPE VALLEY	LOS ANGELES CHILD GUIDANCE	UNITED AMERICAN INDIAN INVOLVEMENT
CHILDRENS HOSPITAL OF LOS ANGELES	MCKINLEY CHILDRENS CENTER	VIP COMMUNITY MENTAL HEALTH CTR INC
CHILDRENS INSTITUTE INC	MENTAL HEALTH AMERICA OF LA	VISTA DEL MAR CHILD AND FAMILY SVC
COUNSELING AND RESEARCH ASSC INC	OLIVE CREST	L.A. COUNTY DMH
COUNSELING4KIDS	OPTIMIST YOUTH HOMES DBA BOYS HOME	EDMUND D. EDELMAN WESTSIDE
CRITTENTON SERV FOR CHILD AND FAMILIES	PACIFIC ASIAN COUN SVS	WEST CENTRAL FAMILY MHS
D VEAL FAMILY AND YOUTH SVC	PACIFIC CLINICS	NORTHEAST MENTAL HEALTH CENTER
DAVID AND MARGARET HOME INC	PACIFIC LODGE YOUTH SERVICES	ARCADIA MHS
DIDI HIRSCH PSYCHIATRIC SVC	PASADENA UNIFIED SCHOOL DISTRICT	LONG BEACH CHILD ADOLESCENT CLINIC
DIGNITY HEALTH DBA CAL HOSP MED CTR	PENNY LANE CENTERS	LONG BEACH MHS ADULT CLINIC
DREW CHILD DEVELOPMENT CORPORATION	PHOENIX HOUSE OF LOS ANGELES INC	SAN PEDRO MENTAL HEALTH
EISNER PEDIATRIC FAMILY CTR	PROVIDENCE COMMUNITY SERVICES	RIO HONDO COMMUNITY MHC
EL CENTRO DE AMISTAD INC	ROSEMARY CHILDRENS SERVICE	SOUTH BAY MHS
EL CENTRO DEL PUEBLO INC	SAINT JOHNS HEALTH CENTER	COMPTON FAMILY MHS
ENKI HEALTH AND RESEARCH SYSTEMS I	SAN FERNANDO VALLEY COMMUNITY MHC	HARBOR-UCLA MEDICA
ETTIE LEE HOMES, INC.	SAN GABRIEL CHILDRENS CENTER INC	AUGUSTUS F HAWKINS FAMILY
EXCEPTIONAL CHILDRENS FOUNDATION	SHIELDS FOR FAMILIES	DOWNTOWN MENTAL HEALTH
FAMILIES FIRST INC	SOCIAL MODEL RECOVERY SYSTEMS	COASTAL API FAMILY MHC
FIVE ACRES	SOUTH CENTRAL HEALTH AND REHAB PROG	LONG BEACH API FAMILY MHC
FOOTHILL FAMILY SERVICE	SPECIAL SERVICE FOR GROUPS	AMERICAN INDIAN COUNSELING
FOR THE CHILD	SPIRITT FAMILY SERVICES	TIES FOR FAMILIES
HAMBURGER HOME DBA AVIVA CENTER	ST ANNES	JUV JUS TRANSITION AFTERCARE
HATHAWAY SYCAMORES CHILD FAM SVC	ST JOSEPH CENTER	

Agencies submitting outcomes that are not approved to provide Seeking Safety by PEI Administration:

ASIAN AMERICAN DRUG ABUSE PROGRAM	L.A. COUNTY DMH
VENICE FAMILY CLINIC	ANTELOPE VALLEY MHS
KEDREN COMMUNITY MENTAL HEALTH CTR	HOLLYWOOD MHC

# of Clients Claimed to Practice	# of Clients entered into PEI OMA	# of Tx cycles in PEI OMA	Clients with Multiple Tx Cycles	Clients Completing Tx	Clients Dropping-Out of Tx	Clients Still in Tx
21464	58.82%	13605	6.89%	26.56%	40.04%	33.40%
n=	12626	n=	870	3614	5447	4544

Note 1: Clients claimed was based on Seeking Safety being selected as the EBP in a PEI Plan and having ≥ 2 core services claimed to the practice.

Note 2: Number of clients Completing Tx or Dropping-Out of Tx was determined by whether the EBP was said to be completed (e.g. answered “yes” or “no”) in the PEI OMA.

Total Number of Clients	Age	Gender			Ethnicity					Primary Language		
	Average	Female	Male	Unknown	African-American	Asian / Pacific Islander	Caucasian	Hispanic / Latino	Other	English	Spanish	Other
12626	22	50.78%	49.16%	0.06%	19.82%	2.30%	11.88%	59.98%	6.03%	76.71%	20.43%	2.86%
n=		6412	6207	7	2502	290	1500	7573	761	9686	2579	361

Note1: Age is calculated at the date of the first EBP.

Note2: Percentages may not total 100 due to missing data and/or rounding.

Total Treatment Cycles	Depressive Disorder NOS	Post-Traumatic Stress Disorder	Oppositional Defiant Disorder	Mood Disorder NOS	Dysthymic Disorder	Other
13605	10.20%	9.59%	9.39%	8.42%	6.56%	55.84%
n=	1388	1305	1277	1145	893	7597

*See breakout of “Other” on the supplemental information page.

Table 4. Program Process Data - Clients Who Entered Seeking Safety			
Outcome Measures Administered	Pre-Test with Scores	Post-test with Scores	Clients Who Completed both a Pre and Post Measure with Scores
UCLA PTSD-RI - Parent n= Ackn=	34.45%	16.30%	7.39%
	3120	964	669
	9057	5914	9057
UCLA PTSD-RI - Child/Adolescent n= Ackn=	63.82%	29.09%	14.97%
	6157	1809	1444
	9647	6218	9647
UCLA PTSD-RI - Adult n= Ackn=	62.68%	27.11%	12.04%
	1958	456	376
	3124	1682	3124
Youth Outcome Questionnaire - 2.01 (Parent) n= Ackn=	46.68%	21.78%	10.32%
	4021	1216	889
	8614	5583	8614
Youth Outcome Questionnaire – Self Report – 2.0 (YOQ-SR) n= Ackn=	74.65%	35.71%	18.64%
	6892	2099	1721
	9232	5878	9232
Outcome Questionnaire - 45.2 n= Ackn=	76.84%	37.04%	16.02%
	2853	717	595
	3713	1936	3713

Note 1: Number of acknowledged measures (Ackn=) is determined by the number of required measures that receive a score or an unable to collect reason code.

Note 2: The % indicated for Pre-test with scores, Post-test with scores, and both a Pre- and Post-test with scores is calculated by dividing the (n=#) by the number acknowledged (Ackn=#) in the PEI OMA system for each measure. The number acknowledged (Ackn=#) for those with Pre and Post scores is an estimate based on the greatest number of matches that could be expected given the number of Pre scores acknowledged.

Table 5a. Top Reasons Given for "Unable to Collect"

UCLA PTSD-RI - Parent	Total Pre 5938	Parent/care provider unavailable	Administration date exceeds acceptable range	Outcome measure unavailable	Parent/care provider refused	Invalid outcome measure	Other Reasons
	Percent	57.02%	14.38%	8.03%	6.89%	4.51%	9.16%
	n	3386	854	477	409	268	544
	Total Post 4950	Parent/care provider unavailable	Premature termination	Lost contact with parent/care provider	Parent/care provider refused	Administration date exceeds acceptable range	Other Reasons
	Percent	51.27%	19.88%	6.67%	5.88%	5.64%	10.67%
	n	2538	984	330	291	279	528

Table 5b. Top Reasons Given for "Unable to Collect"

UCLA PTSD-RI - Child/Adolescent	Total Pre 3490	Administration date exceeds acceptable range	Client refused	Client unavailable	Outcome measure unavailable	Clinician not trained in outcome measure	Other Reasons
	Percent	26.99%	23.55%	14.47%	13.78%	7.59%	13.61%
	n	942	822	505	481	265	475
	Total Post 4409	Premature termination	Client unavailable	Client refused	Lost contact with client	Administration date exceeds acceptable range	Other Reasons
	Percent	28.62%	28.03%	13.15%	10.34%	8.30%	11.54%
	n	1262	1236	580	456	366	509

Table 5c. Top Reasons Given for "Unable to Collect"

UCLA PTSD-RI - Adult	Total Pre 1166	Not available in primary language	Clinician not trained in outcome measure	Outcome measure unavailable	Administration date exceeds acceptable range	Client refused	Other Reasons
	Percent	35.08%	16.64%	15.69%	9.69%	9.18%	13.72%
	n	409	194	183	113	107	160
	Total Post 1226	Client unavailable	Not available in primary language	Lost contact with client	Premature termination	Client refused	Other Reasons
	Percent	20.88%	19.66%	17.05%	16.15%	10.93%	15.33%
	n	256	241	209	198	134	188

Table 5d. Top Reasons Given for "Unable to Collect"							
Youth Outcome Questionnaire - 2.01 (Parent)	Total Pre 4593	Parent/care provider unavailable	Administration date exceeds acceptable range	Parent/care provider refused	Outcome measure unavailable	Clinician not trained in outcome measure	Other Reasons
	Percent	60.85%	16.92%	6.36%	6.21%	3.11%	6.55%
	n	2795	777	292	285	143	301
	Total Post 4367	Parent/care provider unavailable	Premature termination	Lost contact with parent/care provider	Administration date exceeds acceptable range	Parent/care provider refused	Other Reasons
	Percent	52.87%	21.30%	6.76%	6.00%	5.70%	7.37%
	n	2309	930	295	262	249	322

Table 5e. Top Reasons Given for "Unable to Collect"							
Youth Outcome Questionnaire Self Report – 2.0 (YOQ-SR)	Total Pre 2340	Administration date exceeds acceptable range	Client refused	Outcome measure unavailable	Client unavailable	Clinician not trained in outcome measure	Other Reasons
	Percent	34.57%	22.31%	13.59%	12.44%	6.41%	10.68%
	n	809	522	318	291	150	250
	Total Post 3779	Premature termination	Client unavailable	Client refused	Lost contact with client	Administration date exceeds acceptable range	Other Reasons
	Percent	31.89%	29.69%	10.98%	10.74%	8.10%	8.60%
	n	1205	1122	415	406	306	325

Table 5f. Top Reasons Given for "Unable to Collect"							
Outcome Questionnaire – 45.2	Total Pre 860	Outcome measure unavailable	Administration date exceeds acceptable range	Not available in primary language	Client refused	Clinician not trained in outcome measure	Other Reasons
	Percent	22.09%	17.91%	14.65%	11.98%	11.40%	21.98%
	n	190	154	126	103	98	189
	Total Post 1219	Client unavailable	Lost contact with client	Premature termination	Client refused	Not available in primary language	Other Reasons
	Percent	26.83%	21.00%	20.84%	11.81%	6.56%	12.96%
	n	327	256	254	144	80	158

Table 6. Service Delivery Data – Clients Who Completed Seeking Safety						
Total Treatment Cycles 3614	Average Length of Treatment in Weeks	Range of Treatment Weeks		Average Number of Sessions	Range of Sessions	
	32	Min 0	Max 209	35	Min 1	Max 1115

Table 7. Outcome Data[±] – Clients who Completed Seeking Safety					
		Percent Improvement from Pre to Post	Percent of Clients Showing Reliable Change* from Pre-SS to Post-SS		
			Positive Change	No change	Negative Change
UCLA PTSD-RI - Parent	TOTAL	34.43% (n=475)	24.42%	70.95%	4.63%
			116	337	22
UCLA PTSD-RI - Child/Adolescent	TOTAL	28.56% (n=1082)	26.34%	67.10%	6.56%
			285	726	71
UCLA PTSD-RI - Adult	TOTAL	26.37% (n=303)	32.34%	64.36%	3.30%
			98	195	10
Youth Outcome Questionnaire (YOQ) - 2.01 (Parent)	TOTAL	35.30% (n=606)	55.94%	32.18%	11.88%
			339	195	72
Youth Outcome Questionnaire - Self Report - 2.0 (YOQ-SR)	TOTAL	31.60% (n=1329)	43.57%	46.20%	10.23%
			579	614	136
Outcome Questionnaire – 45.2	TOTAL	29.85% (n=486)	54.12%	40.33%	5.56%
			263	196	27

[±]Please see Appendix A. for a description of the Seeking Safety outcome measures and the outcome indicators (percent improvement in average scores; and, percent of clients showing reliable change).

Note1: Possible PTSD-RI range from 0-68, with a clinical cutpoint of 38.



Note2: Possible YOQ Total Scores range from -16-240, with a clinical cutpoint of 46.

Note3: Possible YOQ-SR Total Scores range from -16-240, with a clinical cutpoint of 47.



Note4: Possible OQ Total Scores range from 0-180, with a clinical cutpoint of 64.

Note5: Positive Change indicates that the scores decreased from the pre to the post measures.



UCLA PTSD RI – Parent
(N=475)





UCLA PTSD RI – Child/Adolescent
(N=1082)





UCLA PTSD-RI – Adult
(N=303)





Youth Outcome Questionnaire (YOQ) – 2.01 (Parent)
(N=606)



Youth Outcome Questionnaire – Self Report – 2.0 (YOQ-SR)
(N=1329)



Outcome Questionnaire-45.2 (N=486)



Appendix

Youth Outcomes Questionnaires (YOQ and YOQ-SR)

The Youth Outcome Questionnaire is a 64-item parent-report that assesses global distress in a child's/adolescent's life from 4-17 years of age. The YOQ-SR is the Self-report version of the YOQ and is completed by the child/adolescent. Total scores on both measures can range from -16 to 240. Total scores of 46 or higher are most similar to a clinical population on the YOQ. A total score of 47 is most similar to that of a clinical population on the YOQ-SR.

Outcomes Questionnaires (OQ)

The Outcome Questionnaire is a 45-item self-report that assesses global distress in a client's life from ages 19 and older. Total Scores on this measure can range from 0 to 180, with scores of 64 or higher indicating clinical significance.

Post-Traumatic Stress Disorder Reaction Index (PTSD-RI)

The UCLA Post-Traumatic Stress Disorder Reaction Index (PTSD-RI) is a brief (21 or 22-item measure depending on the version) that measures the frequency of occurrence of post-traumatic stress disorder symptoms during the month prior to the assessment. The Child/Adolescent Version is appropriate for clients age 6-20. The Parent Version is appropriate for clients age 3-18. The Adult Short-form is appropriate for clients age 21+.

Possible Total PTSD Severity Scores range from 0-68; and scores of 38 or higher have the greatest sensitivity and specificity for detecting PTSD.

Reliable Change Index (RCI)

When comparing Pre and Post scores, it is very helpful to know whether the change reported represents the real effects of the treatment or errors in the system of measurement. The Reliability of Change Index (RCI) is a statistical way of helping to insure that the change recorded between pre and post assessments exceeds that which would be expected on the basis of measurement error alone. The RCI has been calculated using the Jacobson and Truax (1991) method and indicates when change exceeds that which would be expected on the basis of error at the $p < .05$ probability level. For a more in-depth discussion of Reliability of Change see Jacobson, N. S., & Truax, P. (1991). *Clinical Significance: A statistical approach to defining meaningful change in psychotherapy research. Journal of Consulting and Clinical Psychology*, 59, 12-19. Also see Wise, E. A. (2004). *Methods for analyzing psychotherapy outcomes: A review of clinical significance, reliable change, and recommendations for future directions. Journal of Personality Assessment*, 82(1), 50-59.

The number and percent of clients experiencing positive change, no change and negative change are recorded in table 7. Healthful change in each of the measures cited here means that scores have decreased in value from pre to post test administrations (i.e. recorded a negative change on the RCI). To help avoid confusion, healthful reliable change is presented as positive while unhealthful reliable change is presented as negative change.

The Reliable Change analysis for the PTSD-RI Adult Short-Form used the same Cronbach Alpha score found in the other PTSD-RI versions.

Supplemental Information for Table 3 “Other” Diagnosis Category

DSM-IV Primary Axis Diagnosis - Clients Who Entered Seeking Safety	n	Percent
Disruptive Behavior Disorder NOS	868	11.43%
Major Depressive Disorder, Recurrent, Moderate	712	9.37%
Conduct Disorder, Adolescent-Onset Type	616	8.11%
Attention-Deficit/Hyperactivity Disorder, Combined Type or Hyperactive Impulse Type	449	5.91%
Major Depressive Disorder, Rec., Severe W/O Psychotic Features	435	5.73%
Generalized Anxiety Disorder	377	4.96%
Anxiety Disorder NOS	367	4.83%
Major Depressive Disorder, Single Episode, Moderate	319	4.20%
Bipolar Disorder NOS	271	3.57%
Major Depressive Disorder, Recurrent, Unspecified	203	2.67%
Major Depressive Disorder, Rec., Severe With Psychotic Features	198	2.61%
Adjustment Disorder With Anxiety and Depressed Mood	186	2.45%
Major Depressive Disorder, Single Episode, Unspecified	179	2.36%
Adjustment Disorder With Depressed Mood	156	2.05%
Major Depressive Disorder, Recurrent, Mild	152	2.00%
Major Depressive Disorder, SE, Severe W/O Psychotic Features	149	1.96%
Major Depressive Disorder, Single Episode, Mild	133	1.75%
Psychotic Disorder NOS	125	1.65%
Attention-Deficit/Hyperactivity Disorder, Predominantly Inattentive Type	111	1.46%
Schizoaffective Disorder	108	1.42%
Adjustment Disorder W/Mixed Disturbance Emotion and Conduct	106	1.40%
Conduct Disorder, Childhood-Onset Type	103	1.36%
Major Depressive Disorder, SE, Severe With Psychotic Features	86	1.13%
Attention-Deficit/Hyperactivity Disorder NOS	73	0.96%
Schizophrenia, Paranoid Type	68	0.90%
Adjustment Disorder Unspecified	63	0.83%
Bipolar II Disorder	63	0.83%
Panic Disorder Without Agoraphobia	60	0.79%
Adjustment Disorder With Anxiety	55	0.72%
Adjustment Disorder With Disturbance of Conduct	47	0.62%
Bipolar I Disorder, MRE, Depressed, Unspecified	37	0.49%
Conduct Disorder, Unspecified Onset	34	0.45%
Bipolar I Disorder, MRE, Depressed, Severe W/O Psychotic Features	33	0.43%
Bipolar I Disorder, MRE, Depressed, Severe With Psychotic Features	33	0.43%
Intermittent Explosive Disorder	33	0.43%
Panic Disorder With Agoraphobia	30	0.39%

Diagnosis Deferred (code invalid as Secondary/Axis II Diagnosis)	27	0.36%
Major Depressive Disorder, Recurrent, In Partial Remission	25	0.33%
Bipolar I Disorder, MRE, Mixed, Unspecified	24	0.32%
DSM-IV Primary Axis Diagnosis - Clients Who Entered Seeking Safety	n	Percent
Bipolar I Disorder, MRE, Mixed, Moderate	24	0.32%
Bipolar I Disorder, MRE, Depressed, Moderate	23	0.30%
Bipolar I Disorder, MRE, Mixed, Severe W/O Psychotic Features	23	0.30%
Social Phobia	22	0.29%
Obsessive-Compulsive Disorder	22	0.29%
Bipolar I Disorder, MRE, Mixed, Severe With Psychotic Features	20	0.26%
Bipolar I Disorder, MRE, Hypomanic/Manic Unspecified	19	0.25%
Bipolar I Disorder, MRE, Manic, Severe With Psychotic Features	19	0.25%
No Diagnosis (code invalid as Secondary/Axis III Diagnosis)	19	0.25%
Bipolar I Disorder, MRE, Manic, Moderate	16	0.21%
Bipolar I Disorder, MRE, Manic, Severe W/O Psychotic Features	15	0.20%
Bipolar I Disorder, MRE, Unspecified	14	0.18%
Mood Disorder Due to ...(Indicate Medical Condition)	14	0.18%
Bipolar I Disorder, MRE, Depressed, Mild	14	0.18%
Acute Stress Disorder	12	0.16%
Schizophrenia, Undifferentiated Type	12	0.16%
Cannabis Abuse	11	0.14%
Schizophrenia, Disorganized Type	11	0.14%
Cyclothymic Disorder	11	0.14%
Reactive Attachment Disorder of Infancy or Early Childhood	10	0.13%
Impulse-Control Disorder NOS	10	0.13%
Separation Anxiety Disorder	9	0.12%
Major Depressive Disorder, Single Episode, In Partial Remission	9	0.12%
Disorder of Infancy, Childhood, or Adolescence NOS	7	0.09%
Pervasive Developmental Disorder NOS/Asperger's Disorder/Rett's Disorder	7	0.09%
Bipolar I Disorder, Single Manic Episode, Unspecified	6	0.08%
Eating Disorder NOS	6	0.08%
Major Depressive Disorder, Recurrent, In Full Remission	6	0.08%
Bereavement	5	0.07%
Delusional Disorder	5	0.07%
Amphetamine Dependence	5	0.07%
Polysubstance Dependence	4	0.05%
Bipolar I Disorder, MRE, Depressed, In Partial Remission	4	0.05%
Drug Induced Mood Disorder	4	0.05%
Drug Induced Psychotic Disorder, With Delusions	3	0.04%
Gender Identity Disorder in Adolescents or Adults	3	0.04%
Alcohol-Induced Anxiety. Mood, Sleep Disorders/Sexual Dysfunction	3	0.04%
Bipolar I Disorder, MRE, Mixed, Mild	3	0.04%
Drug Induced Psychotic Disorder, With Hallucinations	3	0.04%
Bipolar I Disorder, MRE Manic, In Full Remission	2	0.03%

Trichotillomania	2	0.03%
Identity Problem	2	0.03%
Bipolar I Disorder, MRE, Manic, In Partial Remission	2	0.03%
DSM-IV Primary Axis Diagnosis - Clients Who Entered Seeking Safety	n	Percent
Somatoform Disorder NOS and Undifferentiated Somatoform Disorder	2	0.03%
Bulimia Nervosa	2	0.03%
Anxiety Disorder Due to General Medical Condition	2	0.03%
Religious or Spiritual Problem/Phase of Life Problem	2	0.03%
Bipolar I, SME, Severe W/O Psychotic Features	2	0.03%
Anorexia Nervosa	2	0.03%
Major Depressive Disorder, Single Episode, In Full Remission	2	0.03%
Bipolar I Disorder, Single Manic Episode, Moderate	2	0.03%
Bipolar I, SME, Severe With Psychotic Features	2	0.03%
Schizophrenia, Residual Type	2	0.03%
Schizophreniform Disorder	2	0.03%
Opioid Dependence	2	0.03%
Alcohol Dependence	1	0.01%
Drug Induced Delirium	1	0.01%
Dementia due to Medical Disorder/HIV Disease	1	0.01%
Adult Antisocial Behavior	1	0.01%
Conversion Disorder	1	0.01%
Agoraphobia Without History of Panic Disorder	1	0.01%
Pain Disorder Associated w/Psychological and Medical	1	0.01%
Brief Psychotic Disorder	1	0.01%
Dissociative Identity Disorder	1	0.01%
Cannabis Dependence	1	0.01%
Drug Induced Disorder NEC	1	0.01%
Drug Related Disorder NOS	1	0.01%
Psychotic Disorder Due to ...(Indicate Medical Condition) with Delusions	1	0.01%
Parent-Child Relational Problem	1	0.01%
Dyspareunia (Not Due to a Medical Condition)	1	0.01%
Bipolar I Disorder, MRE, Depressed, In Full Remission	1	0.01%
Unspecified Mental Disorder (Nonpsychotic)	1	0.01%
Catatonic Disorder Due to General Medical Condition	1	0.01%
Total	7596	100.00%